

Pi/K Ratio Results

Andre Lebedev
MIPP Software Meeting
May 3, 2007



Approach to Analysis

- Data – MC comparison is not perfect, but looks reasonable
- Compute particle ID variable from mass squared
 - 0=pion
 - 1=proton
 - puts kaon at 0.265
- Extract the occupancy fraction of true pi/K in each histogram
- Apply this fraction to data




Errors

- Statistical
- Correction from MC-reco/MC-ideal comparison
 - Statistical + 0.3 of correction
- Trigger efficiency
- Target out subtraction
- Background modeling
 - Vary (1 – occupancy) by +/-10%



Data-MC positives


MC – yellow shade
Data – red line






Data-MC negatives

MC – yellow shade
Data – red line






Occupancy Positives







Occupancy Negatives






MC Correction and Error






Background Removal






Trigger Efficiency


Trigger efficiency: 1% beryllium, 459 interactions




Trigger efficiency: 1% bismuth, 494 interactions




Trigger efficiency: 2% carbon, 295 interactions




Trigger efficiency: all targets combined




Effect of trigger efficiency on K^+/π^+



Effect of trigger efficiency on K^-/K^+




Effect of trigger efficiency on K^-/π^-






Target out subtraction



Result vs Fluka





What now?

- First analysis result is complete
 - Significant errors from MC/data comparison – need to tune and better understand MC
- I need to finish writing by tomorrow afternoon
 - Will post the thesis to doc-db, comments are welcome in the next 2 weeks